

Double cosets for $SU(2) \times \cdots \times SU(2)$ and outer automorphisms of free groups

YURI A. NERETIN¹

Consider the space of double cosets of the product of n copies of $SU(2)$ with respect to the diagonal subgroup. We get a parametrization of this space, the radial part of the Haar measure, and explicit formulas for the actions of the group of outer automorphisms of the free group F_{n-1} and of the braid group of $n - 1$ strings.

1 Introduction

1.1. The group $SU(2)$. Denote by $SU(2)$ the group of unitary 2×2 -matrices with determinant = 2. A matrix $g \in SU(2)$ has the form

$$g = \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix}, \quad \text{where } |a|^2 + |b|^2 = 1.$$

Therefore, we can identify the manifold $SU(2)$ with the 3-dimensional sphere S^3 in $\mathbb{C}^2 \subset \mathbb{R}^4$.

SS

1.2. Double cosets. Denote by $SU(2)$ the group of unitary 2×2 matrices with determinant = 1. Denote by $G(n)$ the product of n copies of $SU(2)$. Elements of $G(n)$ are n -tuples

$$(g_1, g_2, \dots, g_n), \quad \text{where } g_j \in SU(2). \quad (1.1)$$

Denote by $K = K(n) \simeq SU(2)$ the diagonal subgroup in $G(n)$; elements of K have the form

$$(h, \dots, h), \quad \text{where } h \in SU(2).$$

The object of the paper is the space of double cosets

$$\Pi(n) := K \backslash G/K.$$

In other words, we consider n -tuples (1.1) up to the equivalence

$$(g_1, g_2, \dots, g_n) \sim (hg_1q, \dots, hg_nq), \quad \text{where } h, q \in SU(2).$$

1.3. Conjugacy classes.

Observation 1.1 *There is a canonical one-to-one correspondence between $\Pi(n)$ and the space of conjugacy classes of $G(n - 1)$ with respect to the subgroup $K(n - 1)$.*

¹Supported by the grant FWF, project P19064, by RosAtom, and grants NWO.047.017.015, JSPS-RFBR-07.01.91209.

Indeed,

$$(g_1, g_2, \dots, g_n) \sim (1, g_1^{-1}g_2, \dots, g_1^{-1}g_n)$$

Next,

$$(1, r_1, \dots, r_{n-1}) \sim (1, hr_1h^{-1}, \dots, hr_{n-1}h^{-1}).$$

1.4. Closed polygonal curves on the sphere. Consider the 3-dimensional sphere S^3 endowed with the usual angular distance $d(\cdot, \cdot)$. Fix positive numbers $\theta_1, \dots, \theta_{n-1}$. Consider a closed polygonal curve $A_1A_2 \dots A_{n-1}A_1$ in S^3 such that $d(A_j, A_{j+1}) = \theta_j$, $d(A_{n-1}, A_1) = \theta_{n-1}$. Denote by $\mathcal{X}(\theta)$ the set of all such curves defined up to proper rotations of the sphere.

Observation 1.2 *There is a one-to-one correspondence between $\mathcal{X}(\theta)$ and the set of $(n-1)$ -tuples*

$$(1, r_1, \dots, r_{n-1}), \quad \text{where } r_j \in \text{SU}(2) \quad (1.2)$$

defined up to a simultaneous conjugation and satisfying the conditions:

- *the eigenvalues of r_k are $e^{\pm i\theta_k}$;*
- $r_1r_2 \dots r_{n-1} = 1$.

Indeed, $\text{SU}(2)$ can be considered as a 3-dimensional sphere. To a tuple (1.2), we assign the polygonal curve

$$1, r_1, r_1r_2, \dots, r_1r_2 \dots r_{n-1} = 1.$$

The space $\mathcal{X}(\theta)$ (and its analog for $\mathbb{R}^3$ and the Lobachevsky 3-space) became a subject of investigations after Klyachko's work [12], see e.g. [11], [?]. Relations of the present work with this literature is not quite clear for the author. Some other related works are Fock, Rosly[7], Fock [6], Goldman [9], Dynnikov [5].

1.5. Spectral forms. For a point of $\Pi(n)$, we write the *spectral form*

$$Q(\lambda) := \det\left(\sum_j \lambda_j g_j\right) =: \sum_{i,j} s_{ij} \lambda_i \lambda_j.$$

We describe the set $\Xi(n)$ of possible spectral forms. Namely, they satisfy the conditions:

- $Q(\lambda) \geq 0$;
- $\text{rk } Q \leq 4$;
- $s_{jj} = 1$.

If $\text{rk } Q = 4$, its preimage $\in \Pi(n)$ is a two-point set; we have a branching along the surface $\text{rk } Q = 3$.

Note, that points of the surface $\text{rk } Q = 3$ corresponds to smooth points of the quotient space $\Pi(n) = K \setminus G/K$; the singular locus of $\Pi(n)$ corresponds to the surface $\text{rk } Q \leq 2$.

1.6. Radial part of Haar measure. The group $G(n)$ is endowed with the Haar measure. We consider its pushforward to the space $\Xi(n)$. For $n = 3$

we get the usual Lebesgue measure $ds = ds_{12} ds_{13} ds_{23}$ on $\Xi(3)$, see [14]. For $n = 4$ the measure is given by

$$\det(Q(s))^{-1/2} ds,$$

where ds is the Lebesgue measure. For $n \geq 5$ the description of the measure is given in Theorem 3.5.

1.7. The group $\text{Out}(F_k)$, (see [2], [1]). Consider the free group F_k with k generators $c_1, \dots, c_k$. Denote by $\text{Aut}(F_k)$ the group of automorphisms of F_k . Each automorphism $\varkappa$ is determined by images of the generators:

$$c_j \mapsto \varkappa(c_j) = c_{j_1}^{\varepsilon_{j_1}} c_{j_2}^{\varepsilon_{j_2}} \dots, \quad (1.3)$$

where $\varepsilon_j = \pm 1$. Certainly, the collections $\{\varkappa(c_j)\}$ are not arbitrary (generally, a formula of the type (1.3) determines a non-surjective and non-injective map $F_k \rightarrow F_k$).

By the Nielsen theorem (see [13]), the group $\text{Aut}(F_k)$ is generated by the following transformations of the set of generators:

a) permutations of generators;

b) the map

$$c_1 \mapsto c_1^{-1}, \quad c_2 \mapsto c_2, \quad c_3 \mapsto c_3, \dots;$$

c) the map

$$c_1 \mapsto c_1, \quad c_2 \mapsto c_1 c_2, \quad c_3 \mapsto c_3, \quad c_4 \mapsto c_4, \dots$$

The group F_k acts on itself by interior automorphisms, it is a normal subgroup in $\text{Aut}(F_k)$. We denote by

$$\text{Out}(F_k) := \text{Aut}(F_k)/F_k$$

the group of outer automorphisms of the free group.

1.8. The action of $\text{Out}(F_{n-1})$ on $\Pi(n)$. For a transformation (1.3) we write the following transformation of $\tilde{\Pi}_n$:

$$\tilde{r}_j = r_{j_1}^{\varepsilon_{j_1}} c_{j_2}^{\varepsilon_{j_2}} \dots$$

In Section 4 we obtain explicit formulas for the Nielsen generators.

1.9. Braid groups. See an introduction in [10], and [3], [4], [8]. Denote by Br_k the Artin braid group. It has generators $\sigma_1, \dots, \sigma_{k-1}$ and relations

$$\sigma_j \sigma_{j+1} \sigma_j = \sigma_{j+1} \sigma_j \sigma_{j+1} \quad (1.4)$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i \quad \text{if } |i - j| > 1. \quad (1.5)$$

There is the following Artin embedding $\text{Br}_k \mapsto \text{Aut}(F_k)$ (see [10]). The element h_j corresponds to the transformation

$$c_j \mapsto c_j c_{j+1} c_j^{-1}, \quad c_{j+1} \mapsto c_j,$$

other generators are fixed.

There is a characterization (the Artin theorem) of the image of Br_k in $\text{Aut}(F_k)$. Namely, $\varkappa \in \text{Out}(F_k)$ is contained in Br_k if

1) $\varkappa$ sends each generator c_j to

$$A_j^{-1} c_{\xi(j)} A_j$$

where $A_j \in F_k$ and ξ is a permutation of generators.

2) $\varkappa$ sends $c_1 \dots c_k$ to itself.

In particular, we get the map $\text{Br}_k \rightarrow \text{Out}(F_k)$. It is not injective (see, e.g., [8]), the kernel is generated by

$$\left((\sigma_1 \sigma_2 \dots \sigma_{k-1}) (\sigma_1 \sigma_2 \dots \sigma_{k-2}) \dots \sigma_1 \right)^2.$$

In Section 4 we get explicit formulas for the action of generators of the braid group in the terms of spectral forms.

1.10. Pure braids. The relations

$$\sigma_j^2 = 1$$

together with (??)–(??) determine the symmetric group S_n . Therefore we get the homomorphism of $\text{Br}_n \rightarrow S_n$. The kernel is called the *group of pure braids*. The group of pure braids acts on F_n by transformations of the form

$$c_j \mapsto A_j c_j A_j^{-1}$$

recall that $c_1 \dots c_n \mapsto_1 \dots c_n$. This implies the following observation

Observation 1.3 *The group of pure braids acts on the space $\mathcal{X}(\theta)$ of closed polygonal curves.*

1.11. The structure of the paper. In Section 2, we get the characterization of spectral forms. Section 3 contains evaluation of the radial part of the Haar measure. In Section 4, we write out actions of the the groups $\text{Out}(F_{n-1})$ and Br_{n-1} .

I am grateful to R. S. Ismagilov, V. A. Fock, A. A. Rosly, L. G. Rybnikov, and I. A. Dynnikov for discussion of this topic.

2 Spectral forms

2.1. Spectral forms and the map ζ . For any element of $\Pi(n)$ we write out the quadratic form

$$Q(\lambda_1, \dots, \lambda_n) = \det\left(\sum \lambda_j g_j\right) = \det\left(\sum \lambda_j \begin{pmatrix} a_j & b_j \\ -\bar{b}_j & \bar{a}_j \end{pmatrix}\right)$$

We denote by ζ the map from $\Pi(n)$ to the space of quadratic forms.

Proposition 2.1 *a) We get a well-defined map from the space $\Pi(n)$ to the space of quadratic forms.*

b) Coefficients of Q are real, coefficients in the front of λ_j^2 are 1.

c) Q is positive semidefinite.

d) The rank of Q is ≤ 4 .

PROOF. Indeed, for real λ ,

$$\begin{aligned} Q(\lambda) &= \left(\sum \lambda_j a_j \right) \left(\sum \lambda_j \bar{a}_j \right) + \left(\sum \lambda_j b_j \right) \left(\sum \lambda_j \bar{b}_j \right) = \\ &= \left| \sum \lambda_j a_j \right|^2 + \left| \sum \lambda_j b_j \right|^2 = \sum \lambda_j^2 + 2 \sum_{i < j} \operatorname{Re}(a_i \bar{a}_j + b_i \bar{b}_j) \end{aligned}$$

and all the statements become obvious. $\square$

2.2. A description of $\Pi(n)$. Denote by $\Xi = \Xi(n)$ the set of all quadratic forms Q satisfying the conditions of the previous statement.

Obviously,

$$\zeta(g_1^t, \dots, g_n^t) = \zeta(g_1, \dots, g_n),$$

where ^{t} denotes the transposed matrix.

Theorem 2.2 *a) The map $\zeta : \Pi(n) \rightarrow \Xi(n)$ is surjective.*

b) The ζ -preimage of a point $Q \in \Xi$ consists of two points if $\operatorname{rk} Q = 4$ and of one point if $\operatorname{rk} Q \leq 3$.

c) Moreover, $\operatorname{rk} Q \leq 3$ iff $(g_1^t, \dots, g_n^t)$ and $(g_1, \dots, g_n)$ represent one point of Π .

PROOF. For a positive semi-definite quadratic form

$$Q(\lambda) = \sum_{kl} s_{kl} \lambda_k \lambda_l$$

on $\mathbb{R}^n$ there is a collection (a configuration) of vectors v_j in a Euclidean space such that

$$\langle v_k, v_j \rangle = s_{kj}$$

Since $\operatorname{rk} Q \leq 4$, this configuration can be realized in $\mathbb{R}^4$. Since $s_{jj} = 1$, these vectors lie on the unit sphere.

Moreover such a configuration $v_j \in \mathbb{R}^4$ is unique up to the action of the orthogonal group $O(4)$.

Recall that $SU(2)$ can be identified with the 3-dimensional sphere S^3 , then s_{ij} are inner products of points of the sphere. Recall that $SO(4) \simeq SU(2) \times SU(2)/\{\pm 1\}$. In other words, proper isometries of the sphere S^3 correspond to the left-right action of $SU(2) \times SU(2)$ on $SU(2)$ (see, e.g. [15]).

If $\operatorname{rk} Q = 4$, then v_j are not contained in a 3-dimensional hyperplane. Therefore an improper isometry of the sphere gives a non-equivalent configuration in $\Pi(n)$.

If $\operatorname{rk} Q \leq 3$, then the point configuration v_j is contained in a hyperplane. The reflection with respect to the hyperplane fix this configuration. $\square$

3 The radial part of the Haar measure

3.1. A reduction. As we noted above, any element of the double coset space $\Pi(n)$ can be reduced to the form $(1, g_2, \dots, g_n)$ and g_j are determined up to a simultaneous conjugation.

Proposition 3.1 *Each element of $\Pi(n)$ has a representative of the form*

$$\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} e^{i\varphi} & 0 \\ 0 & e^{-i\varphi} \end{pmatrix}, \begin{pmatrix} a_2 & b_2 \\ -\bar{b}_2 & \bar{a}_2 \end{pmatrix}, \dots, \begin{pmatrix} a_{n-1} & b_{n-1} \\ -\bar{b}_{n-1} & \bar{a}_{n-1} \end{pmatrix} \right\},$$

where $b_2 \geq 0$ and $0 \leq \varphi \leq \pi$. (3.1)

For an element in a general position such a representative is unique.

Indeed, after a reduction of g_2 to a diagonal form, we can conjugate our tuple by diagonal matrices.

3.2. The Haar measure on $SU(2)$. We can regard the group $SU(2)$ as the unit sphere in the Euclidean space $\mathbb{C}^2$. The Haar measure on $SU(2)$ is the usual surface Lebesgue measure on the sphere. We denote this measure by dg .

Recall the following simple facts.

Proposition 3.2 *a) The image of the Haar measure under the map $\begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \mapsto a$ is the Lebesgue measure $da d\bar{a}$ on the circle $|a| \leq 1$.*

b) Represent b in the form $b = \rho e^{i\theta}$. Then the image of the Lebesgue measure under the map $\begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \mapsto (a, \theta)$ is

$$\text{const} \cdot d\theta da d\bar{a}.$$

c) Consider the map taking a matrix g to its collection of eigenvalues $e^{i\varphi}, e^{-i\varphi}$, where $0 \leq \varphi \leq \pi$. The image of the Haar measure under the map $g \mapsto \varphi$ is $\sin^2 \varphi d\varphi$.

Corollary 3.3 *The pushforward of the Haar measure in the coordinates (3.1) is*

$$\sin^2 \varphi d\varphi da_2 d\bar{a}_2 dg_3 \dots dg_n$$

3.3. Coordinates. n-tuples of matrices. To be definite, take $n = 5$,

$$(g_1, g_2, g_3, g_4, g_5) := \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} e^{i\varphi} & 0 \\ 0 & e^{-i\varphi} \end{pmatrix}, \begin{pmatrix} a_1 & b_1 \\ -\bar{b}_1 & \bar{a}_1 \end{pmatrix}, \begin{pmatrix} a_2 & b_2 \\ -\bar{b}_2 & \bar{a}_2 \end{pmatrix}, \begin{pmatrix} a_3 & b_3 \\ -\bar{b}_3 & \bar{a}_3 \end{pmatrix} \right),$$

where $0 \leq \varphi \leq \pi$. We also denote

$$a_1 =: x_1 + iy_1 \quad a_2 =: x_1 + iy_2 \quad a_3 = x_3 + iy_3, \quad (3.2)$$

$$b_1 =: e^{i\theta_1} \sqrt{1 - x_1^2 - y_1^2} \quad b_2 =: e^{i\theta_2} \sqrt{1 - x_2^2 - y_2^2} \quad b_3 =: e^{i\theta_3} \sqrt{1 - x_3^2 - y_3^2}. \quad (3.3)$$

For $(g_1, g_2, g_3, g_4, g_5) \in \Pi(n)$ the numbers

$$\theta_1 - \theta_2, \quad \theta_2 - \theta_3, \quad \theta_1 - \theta_3$$

make sense (but not $\theta_1, \theta_2, \theta_3$ themselves).

3.4. Coordinates. Spectral forms. Consider the spectral form and denote its coefficients in the following way:

$$\begin{aligned} \det \left(\sum \lambda g_1 + \mu g_2 + \nu_1 g_3 + \nu_2 g_4 + \nu_3 g_5 \right) =: \\ =: \lambda^2 + \mu^2 + \nu^2 + 2p\lambda\mu + 2q_1\lambda\nu_1 + 2q_2\lambda\nu_2 + 2q_3\lambda\nu_3 \\ + 2r_1\mu\nu_1 + 2r_2\mu\nu_2 + 2r_3\mu\nu_3 + 2t_{12}\nu_1\nu_2 + 2t_{13}\nu_1\nu_3 + 2t_{23}\nu_2\nu_3. \end{aligned}$$

The matrix of the form is

$$\Delta = \begin{pmatrix} 1 & p & q_1 & q_2 & q_3 \\ p & 1 & r_1 & r_2 & r_3 \\ q_1 & r_1 & 1 & t_{12} & t_{13} \\ q_2 & r_2 & t_{12} & 1 & t_{23} \\ q_3 & r_3 & t_{13} & t_{23} & 1 \end{pmatrix}. \quad (3.4)$$

Then

$$p = \cos \varphi \quad (3.5)$$

$$q_j = x_j \quad (3.6)$$

$$r_j = x_j \cos \varphi + y_j \sin \varphi \quad (3.7)$$

$$t_{ij} = x_i x_j + y_i y_j + \sqrt{1 - x_i^2 - y_i^2} \sqrt{1 - x_j^2 - y_j^2} \cos(\theta_i - \theta_j) \quad (3.8)$$

It is easy to write the inverse map:

$$\varphi = \arccos p \quad (3.9)$$

$$x_j = q_j \quad (3.10)$$

$$y_j = \frac{r_j - q_j p}{\sqrt{1 - p^2}} \quad (3.11)$$

$$\theta_i - \theta_j = \pm \arccos \frac{\det \begin{pmatrix} 1 & p & q_i \\ p & 1 & r_i \\ q_j & r_j & t \end{pmatrix}}{\det^{1/2} \begin{pmatrix} 1 & p & q_i \\ p & 1 & r_i \\ q_i & r_i & 1 \end{pmatrix} \det^{1/2} \begin{pmatrix} 1 & p & q_j \\ p & 1 & r_j \\ q_j & r_j & 1 \end{pmatrix}} \quad (3.12)$$

The last formula requires some calculations. For this reason, we present some intermediate formulas:

$$\begin{aligned} 1 - x_1^2 - y_1^2 &= \frac{\det \begin{pmatrix} 1 & p & q_1 \\ p & 1 & r_1 \\ q_1 & r_1 & 1 \end{pmatrix}}{1 - p^2}, \\ t - x_1 x_2 - y_1 y_2 &= \frac{\det \begin{pmatrix} 1 & p & q_1 \\ p & 1 & r_1 \\ q_2 & r_2 & t \end{pmatrix}}{1 - p^2}. \end{aligned} \quad (3.13)$$

Note, that we can not reconstruct the sign of $\theta_i - \theta_j$ from the formula (3.12). Recall that the substitution

$$\theta_1 \mapsto -\theta_1, \quad \theta_2 \mapsto -\theta_2, \quad \theta_3 \mapsto -\theta_3$$

corresponds to the simultaneous transposition

$$(g_1, g_2, g_3, g_4, g_5) \mapsto (g_1^t, g_2^t, g_3^t, g_4^t, g_5^t).$$

3.5. What happens if we forget t_{23} ? Let we know p , all q_j , all r_j , and t_{12}, t_{13} . Then we can reconstruct φ, x_j, y_j and

$$\cos(\theta_1 - \theta_2), \quad \cos(\theta_1 - \theta_3).$$

Without loss of a generality, we can assume $\theta_1 = 0$. Then we know $\pm\theta_2$ and $\pm\theta_3$ and there are two possible variants for $|\theta_2 - \theta_3|$.

It can be readily checked that there exist $h \in \text{SU}(2)$ such that

$$h^{-1} g_2 h = g_2^t, \quad h^{-1} g_3 h = g_3^t$$

(recall that g_1 is the unit matrix). Then without t_{23} we can not distinguish

$$(g_1, g_2, g_3, g_4, g_5) \quad \text{and} \quad (g_1, g_2, g_3, g_4, h g_5^t h^{-1}). \quad (3.14)$$

3.6. The radial part of the Haar measure. The cases $n = 3, n = 4$.

Theorem 3.4 *a) Let $n = 3$. The pushforward of the Haar measure under the map $\zeta : \Pi(3) \rightarrow \Xi(3)$ is²*

$$\text{const} \cdot dp dq_1 dr_1.$$

b) Let $n = 4$. Then the image of the Haar measure under the map $\zeta : \Pi(4) \rightarrow \Xi(4)$ is

$$\text{const} \cdot \det \begin{pmatrix} 1 & p & q_1 & q_2 \\ p & 1 & r_1 & r_2 \\ q_1 & r_1 & 1 & t_{12} \\ q_2 & r_2 & t_{12} & 1 \end{pmatrix}^{-1/2} dp dq_1 dq_2 dr_1 dr_2 dt_{12}.$$

²See, also [9].

PROOF. Consider the case $n = 4$. The radial part of the Haar measure in the coordinates $\varphi, x_1, y_1, x_2, y_2, \theta$ is given by

$$\text{const} \cdot \sin^2 \varphi d\varphi dx_1 dy_1 dx_2 dy_2 d\theta$$

Next, we must write the Jacobian of the map (3.9)–(3.12). Evidently, the Jacobian is

$$\frac{\partial \varphi}{\partial p} \cdot \frac{\partial \theta}{\partial t},$$

this can be easily evaluated.

3.7. The Haar measure, general case. For an n -tuple $(g_1, \dots, g_n)$ consider its spectral form

$$\det\left(\sum_j \lambda_j g_j\right) =: \sum_j \lambda_j^2 + 2 \sum_{i < j} s_{ij} \lambda_i \lambda_j$$

Theorem 3.5 a) *The coefficients s_{12}, s_{13}, s_{23} are distributed as*

$$\text{const} \cdot ds_{12} ds_{13} ds_{23}$$

$$\text{in the domain } \begin{pmatrix} 1 & s_{12} & s_{13} \\ s_{12} & 1 & s_{23} \\ s_{13} & s_{23} & 1 \end{pmatrix} \geq 0.$$

b) *For fixed s_{12}, s_{13}, s_{23} in a general position, a vector $v_j := (s_{1j} \ s_{2j} \ s_{3j})$ is distributed as*

$$\text{const} \cdot \det(\Delta_j)^{-1/2} ds_{j1} ds_{j2} ds_{j3},$$

where

$$\Delta_j = \begin{pmatrix} 1 & s_{12} & s_{13} & s_{1j} \\ s_{12} & 1 & s_{23} & s_{2j} \\ s_{13} & s_{23} & 1 & s_{3j} \\ s_{1j} & s_{2j} & s_{3j} & 1 \end{pmatrix}.$$

A vector $(s_{1j} \ s_{2j} \ s_{3j})$ ranges in the domain $\Delta_j \geq 0$.

c) *The random variables $v_4, v_5, \dots, v_n$ are independent.*

d) *Let us fix s_{1j}, s_{2j}, s_{3j} for all j . For such a collection in a general position there are 2 equiprobable variants of a choice of s_{4j} . These samplings are independent for $j = 5, 6, \dots$*

e) *Let us fix s_{1j}, s_{2j}, s_{3j} for all j and fix s_{4j} . Then a.s. all other variables s_{ij} are uniquely determined.*

PROOF. The statements a)-b) are a rephrasing of Theorem (3.4).

Next, for fixed g_1, g_2, g_3 the matrices ('random variables') $g_3, g_4, \dots$ are independent. A matrix g_j determines a vector v_j , and g_j is uniquely determined by a vector v_j . This proves c).

Now we write the matrix of a spectral form

$$\begin{pmatrix} 1 & s_{12} & s_{13} & s_{14} & s_{15} & s_{16} & \dots \\ s_{12} & 1 & s_{23} & s_{24} & s_{25} & s_{26} & \dots \\ s_{13} & s_{23} & 1 & s_{34} & s_{35} & s_{36} & \dots \\ s_{14} & s_{24} & s_{34} & 1 & ?_1 & ?_2 & \dots \\ s_{15} & s_{25} & s_{35} & ?_1 & 1 & * & \dots \\ s_{16} & s_{26} & s_{36} & ?_2 & * & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The left upper 5×5 minor is zero. This gives a quadratic equation for s_{45} . Both the solutions are admissible, they correspond to collections of matrices given by (3.14). We repeat the same operation for s_{46} etc.

In the terminology of Subsections 3.4–3.5, the knowledge of the first three rows gives us $\pm\theta_1, \pm\theta_2, \pm\theta_3$ etc. All these choices are equiprobable.

Having all s_{4j} , we get a unique way to complete the matrix to the matrix of rank 4.

4 Actions of $\text{Out}(F_{n-1})$ and braid group

4.1. The action of $\text{Out}(F_{n-1})$ on $\Pi(n)$. We regard $\Pi(n)$ as the set of collections $(1, g_2, \dots, g_n)$ defined up to a simultaneous conjugation.

The Nielsen transformations act on $\Pi[n]$ in the obvious way. We get:

- a) permutations of g_j ;
- b) the transformation $g_2 \mapsto g_2^{-1}$;
- c) the map $(g_2, g_3, g_4, \dots) \rightarrow (g_2, g_2g_3, g_4, \dots)$.

To be definite, take $n = 5$. The action of permutations is obvious.

Proposition 4.1 *The transformation $g_2 \mapsto g_2^{-1}$ corresponds to the map $\Xi(5) \rightarrow \Xi(5)$ given by*

$$\begin{aligned} \tilde{p} &= p; \\ \tilde{q}_j &= q_j; \\ \tilde{t}_{ij} &= t_{ij} \\ \tilde{r}_j &= -r_j + 2pq_j \end{aligned}$$

PROOF. In the notation (3.2)–(3.3), (3.5)–(3.8), we have

$$\tilde{r}_1 = x_1 \cos \varphi - x_1 \sin \varphi = r_1 - 2y_1 \sin \varphi.$$

On the other hand,

$$y_1 \sin \varphi = r_1 - x_1 \cos \varphi = r_1 - pq_1.$$

This completes the calculation. $\square$

Theorem 4.2 *The transformation*

$$(1, g_2, g_3, g_4, g_5) \mapsto (1, g_2, g_2 g_3, g_4, g_5)$$

corresponds to the map $\tilde{\Xi}(5) \rightarrow \tilde{\Xi}(5)$ given by

$$\begin{aligned} \tilde{p} &= p; \\ \tilde{q}_2 &= q_2, \quad \tilde{q}_3 = q_3; \\ \tilde{q}_1 &= -r_1 + 2pq_1; \\ \tilde{r}_2 &= r_2, \quad \tilde{r}_3 = q_3; \\ \tilde{t}_{1j} &= p t_{1j} - q_j r_1 + q_1 r_j \mp \det \begin{pmatrix} 1 & p & q_1 & q_j \\ p & 1 & r_1 & r_j \\ q_1 & r_1 & 1 & t_{1j} \\ q_j & r_j & t_{1j} & 1 \end{pmatrix}^{1/2}, \quad \text{where } j = 2, 3. \\ \tilde{t}_{23} &= t_{23} \end{aligned} \quad (4.1)$$

The group $\text{Out}(F_n)$ acts on $\tilde{\Xi}(5)$ and not on $\Xi(5)$ and the choice of signs $\mp$ requires explanations. They are given below.

PROOF. First, we write the coefficients of the spectral form for the transformed collection. Only the variables q_1, r_1, t_{12}, t_{13} change. We have

$$\begin{aligned} \tilde{r}_1 &= \tilde{x}_1 \cos \varphi + \tilde{y}_1 \sin \varphi = \\ &= (x_1 \cos \varphi - y_1 \sin \varphi) \cos \varphi + (y_1 \cos \varphi + x_1 \sin \varphi) \sin \varphi = x_1 = q_1. \end{aligned}$$

and

$$\tilde{q}_1 = \tilde{x}_1 = x_1 \cos \varphi - y_1 \sin \varphi = q_1 p - (r_1 - q_1 p) = -r_1 + 2q_1 p. \quad (4.2)$$

The evaluation of $\tilde{t}_{1j}$ is heavier,

$$\tilde{t}_{12} = \tilde{x}_1 \tilde{x}_2 + \tilde{y}_1 \tilde{y}_2 + \sqrt{1 - \tilde{x}_1^2 - \tilde{y}_1^2} \sqrt{1 - \tilde{x}_2^2 - \tilde{y}_2^2} \cos(\tilde{\theta}_1 - \tilde{\theta}_2).$$

By construction, $\tilde{x}_2 = x_2, \tilde{y}_2 = y_2, \tilde{\theta}_2 = \theta_2$. Next,

$$\tilde{a}_1 = \tilde{x}_1 + i\tilde{y}_1 = (x_1 + iy_1) e^{i\varphi}$$

and therefore

$$1 - \tilde{x}_1^2 - \tilde{y}_1^2 = 1 - x_1^2 - y_1^2.$$

Also, $\tilde{\theta}_1 = \theta_1 + \varphi$. Denote $\theta := \theta_1 - \theta_2$. Thus,

$$\tilde{t}_{12} = \tilde{x}_1 x_2 + \tilde{y}_1 y_2 + \sqrt{1 - x_1^2 - y_1^2} \sqrt{1 - x_2^2 - y_2^2} (\cos \theta \cos \varphi - \sin \theta \sin \varphi).$$

The variable $\tilde{x}_1$ is evaluated in (4.2),

$$\tilde{y}_1 = x_1 \sin \varphi + y_1 \cos \varphi = \frac{r_1 - q_1 p}{\sqrt{1 - p^2}} \cdot p + q \sqrt{1 - p^2}.$$

We use formula (3.13) for square roots and formula (3.12) for $\cos \theta$. After this, we can evaluate $\sin \theta$,

$$\sin^2 \theta = \frac{(1 - p^2) \cdot \det \begin{pmatrix} 1 & p & q_1 & q_2 \\ p & 1 & r_1 & r_2 \\ q_1 & r_1 & 1 & t_{12} \\ q_2 & r_2 & t_{12} & 1 \end{pmatrix}}{\det \begin{pmatrix} 1 & p & q_1 \\ p & 1 & r_1 \\ q_1 & r_1 & 1 \end{pmatrix} \det \begin{pmatrix} 1 & p & q_2 \\ p & 1 & r_2 \\ q_2 & r_2 & 1 \end{pmatrix}}.$$

After this, we get a unexpectedly long chain of cancelations and get the desired formula. $\square$

CHOICE OF SIGNS. We use formula (3.12) and find

$$\pm(\theta_1 - \theta_2), \quad \pm(\theta_1 - \theta_3), \quad \pm(\theta_2 - \theta_3)$$

These numbers must be consistent, in fact only two variants are possible (this corresponds to a choice of a sheet of the covering map $\tilde{\Xi} \rightarrow \Xi$). Now let we have chosen the signs. Then we take 'minus' in (4.1) if $(\theta_1 - \theta_j) \geq 0$. Otherwise, we take 'plus'.

4.2. The action of the braid group.

Lemma 4.3 *The transformation*

$$(1, g_2, g_3, g_4, g_5) \mapsto (1, g_2 g_3 g_2^{-1}, g_2, g_4, g_5)$$

corresponds to the map $\tilde{\Xi}(5) \rightarrow \tilde{\Xi}(5)$ given by

$$\begin{aligned} \tilde{p} &= p \\ \tilde{q}_k &= q_k, \quad \text{where } k = 1, 2, 3; \\ \tilde{r}_k &= r_k, \quad \text{where } k = 1, 2, 3; \\ \tilde{t}_{1j} &= t_{1j} - 2 \cdot \det \begin{pmatrix} 1 & p & q_1 \\ p & 1 & r_1 \\ q_j & r_j & t_{1j} \end{pmatrix} \mp 2p \cdot \det \begin{pmatrix} 1 & p & q_1 & q_j \\ p & 1 & r_1 & r_j \\ q_1 & r_1 & 1 & t_{1j} \\ q_j & r_j & t_{1j} & 1 \end{pmatrix}^{1/2}, \\ \tilde{t}_{23} &= t_{23}. \end{aligned}$$

PROOF. We evaluate

$$\begin{aligned} \tilde{t}_{12} &= \operatorname{Re}(a_1 \bar{a}_2 + b_1 \bar{b}_2 e^{2i\varphi}) = \\ &= x_1 x_2 + y_1 y_2 + \sqrt{1 - x_1^2 - y_1^2} \sqrt{1 - x_2^2 - y_2^2} (\cos \theta \cos 2\varphi - \sin \theta \sin 2\varphi) \end{aligned}$$

as in the previous proof. $\square$

Now we can write the action of the braid group. To write formulas for generators, we represent the matrix of the spectral form as

$$\begin{pmatrix} 1 & p_1 & p_2 & p_3 & \dots \\ p_1 & 1 & h_{12} & h_{13} & \dots \\ p_2 & h_{12} & 1 & h_{23} & \dots \\ p_3 & h_{13} & h_{23} & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

We also set $h_{ij} := h_{ji}$. Then the formula for a generator σ_k of the braid group is

$$\begin{aligned} \tilde{h}_{ij} &= h_{ij}, & \text{if } i, j \neq k, k+1 \\ \tilde{p}_i &= p_i, & \text{if } i \neq k, k+1 \\ \tilde{p}_k &= p_{k+1} \\ \tilde{p}_{k+1} &= p_k \\ \tilde{h}_{(k+1)j} &= h_{kj}, \end{aligned}$$

and

$$\begin{aligned} \tilde{h}_{kj} &= h_{(k+1)j} - 2 \cdot \det \begin{pmatrix} 1 & p_k & p_{k+1} \\ p_k & 1 & h_{k(k+1)} \\ p_j & h_{kj} & h_{(k+1)j} \end{pmatrix} - \\ &\quad - 2p_k \cdot \det \begin{pmatrix} 1 & p_k & p_{k+1} & p_j \\ p_k & 1 & h_{k(k+1)} & h_{kj} \\ p_{k+1} & h_{k(k+1)} & 1 & h_{(k+1)j} \\ p_j & h_{kj} & h_{(k+1)j} & 1 \end{pmatrix}^{1/2} \end{aligned} \quad (4.3)$$

References

- [1] Bestvina, M. *The topology of $Out(F_n)$* . Proceedings of ICM, Beijing, 2002, v.2, 373–384.
- [2] Culler, M., Vogtmann, K., *Moduli of graphs and automorphisms of free groups*. Inv. Math., 84 (1986), 91–119
- [3] Dehornoy, P., *Braids and self-distributivity*, Birkhauser, 2000
- [4] Dehornoy, P., Dynnikov, I., Rolfsen, D., Weist, B., *Orderings of braids*. Amer. Math. Soc., 2009
- [5] Dynnikov, I. A. *On a Yang–Baxter map and Dehornoy ordering*, Russ. Math. Surv., 57 (2002) 592–594
- [6] Fock, V. V. *Dual Teichmüller space*, Theoret. Math. Phys., 120(1999), 1245–1259

- [7] Fock, V. V.; Rosly, A. A. *Flat connections and polyubles*. Teoret. Mat. Fiz. 95 (1993), no. 2, 228–238; translation in Theoret. and Math. Phys. 95 (1993), no. 2, 526–534
- [8] Garber, D. *Braid group cryptography*, Preprint, available via <http://arXiv.org/abs/0711.3941>
- [9] Goldman, W. M. *An ergodic action of the outer automorphism group of a free group*. Geom. Funct. Anal. 17 (2007), no. 3, 793–805.
- [10] Hansen, V. L. *Braids and coverings: selected topics*. Cambridge University Press, 1989
- [11] Kapovich, M., Millson, John J., Treloar, T. *The symplectic geometry of polygons in hyperbolic 3-space*. Asian J. Math. 4 (2000), no. 1, 123–164.
- [12] Klyachko, A. A. *Spatial polygons and stable configurations of points on a projective line*. Algebraic geometry and applications, Yaroslavl, 1992, 67–84; English transl. in Aspects Math., E25, Vieweg, Braunschweig, 1994
- [13] Lyndon, R. C., Schupp, P. *Combinatorial group theory*, Springer, 1977
- [14] Neretin, Yu. A. *On spherical functions on the group $SU(2) \times SU(2) \times SU(2)$* , Funct. Anal. Appl., to appear.
- [15] Zhelobenko, D. P. *Compact Lie groups and their representations*, American Mathematical Society, Providence, R.I., 1973.

Math.Dept., University of Vienna,
 Nordbergstrasse, 15, Vienna, Austria
 &
 Institute for Theoretical and Experimental Physics,
 Bolshaya Cheremushkinskaya, 25, Moscow 117259, Russia
 &
 Mech.Math. Dept., Moscow State University, Vorob'evy Gory, Moscow
 e-mail: neretin(at) mccme.ru
 URL: www.mat.univie.ac.at/~neretin
wwwth.itep.ru/~neretin